

WHY STUDY INTERSTELLAR MATTER VERY CLOSE TO THE SUN?

P.C. Frisch¹

¹*University of Chicago, Department of Astronomy and Astrophysics, 5640 South Ellis Avenue,
Chicago, IL 60637 USA*

ABSTRACT

Interstellar matter (ISM) sets the boundary conditions of the heliosphere and dominates the interplanetary medium. The heliosphere configuration has varied over recent history, as the Sun emerged from the Local Bubble and entered a turbulent outflow of cloudlets originating from the Scorpius-Centaurus Association direction. Several indicators suggest that the local interstellar magnetic field is weak and parallel to the galactic plane. Observations of the interaction products of the ISM with the heliosphere, such as pickup ions and anomalous cosmic rays, when combined with data on the ISM towards nearby stars, provide unique constraints on the composition and physical properties of nearby gas. These data suggest abundances in nearby ISM are subsolar, and that gas and dust are not well mixed at the solar location.

INTRODUCTION

Interstellar gas and dust within ~ 30 pc of the Sun provide a unique opportunity to evaluate the chemical composition and physical properties of diffuse interstellar clouds in a small region of space. Several motivations for studying local interstellar matter (ISM, LISM) stand out.

1. The ISM sets the boundary conditions of the heliosphere. By analogy, the ISM also sets the boundary conditions for the astrospheres of extrasolar planetary systems. Through the heliosphere-cosmic ray flux (and by analogy the astrosphere-cosmic ray flux) connection, the distribution of stable planetary climates may, in part, depend on the present and historical physical properties of the ISM surrounding each system.
2. Although many details of the chemical composition of the ISM are known, the overall picture is poorly understood because elements are observed only in the gas phase of the ISM, line blending in distant stars confuses the interpretation of data, H^+ is unobserved in low column density sightlines, and the interstellar grain composition, gas-dust mixing and chemical homogeneity are unknown.
3. The configuration of the heliosphere depends on the physical properties of the surrounding ISM. The Sun moves through space, and currently the Sun is embedded in an outflow of material from the Scorpius constellation. As the Sun travels through space, the heliosphere is responsive to “galactic weather” patterns. Also, the plane of the ecliptic is tilted by $\sim 60^\circ$ with respect to the galactic plane, introducing a north/south asymmetry in the heliosphere because the local interstellar magnetic field appears to be parallel to the galactic plane.

4. Turbulence in the ISM, whether macroscopic or microscopic, is poorly understood. Interstellar absorption lines at the velocity of the cloud surrounding the solar system (also known as the Local Interstellar Cloud, LIC) are not observed towards the nearest star, α Cen ($\sim 45^\circ$ from the nose of the heliosphere), suggesting either a turbulence-driven velocity discontinuity, or a physically distinct interstellar cloudlet in this direction. The heliosphere responds differently to the case where velocity discontinuities, versus “empty” space, distinguish cloudlets.
5. Lastly, dust and neutral gas from the LIC enter the solar system, where observations of the interaction products directly sample the chemical composition and ionization levels of the parent ISM. The combination of ISM observations towards nearby stars and *in situ* observations of He, H, pickup ions, and anomalous cosmic rays in the heliosphere provides strong constraints on the physical conditions of the surrounding cloud.

These unique qualities of the ISM close to the Sun offer a space laboratory for studying cosmic processes that is not duplicated elsewhere in the Universe, and are elaborated on below.

BOUNDARY CONDITIONS OF HELIOSPHERE AND ASTROSPHERES

The first spectral observations of the interplanetary Lyman- α glow established that neutral gas at interstellar velocities has penetrated the heliosphere (Adams & Frisch 1977). The heliosphere responds to the relative Sun-cloud velocity and the partial pressure of the constituents of the LIC. As the Sun journeys through the Milky Way Galaxy, the heliosphere configuration varies in response to the properties of the surrounding ISM (Frisch 1993, 1997; Zank & Frisch 1999; Mueller et al. 2001, 2002). Using recent Hipparcos results for the solar apex motion (Dehnen & Binney 1998), the Sun moves through the local standard of rest at ~ 14 parsecs per million years, and has a relative velocity with respect to the LIC of ~ 26 parsecs per million years.

The Sun is currently immersed in an outflow of ISM from the region of the Scorpius-Centaurus Association (see below), and the distance to the outflow surface in the heliosphere tail direction is ~ 0.9 pc. Hence the Sun is likely to have entered this outflow within the past 10^5 years, and possibly within the past $\sim 10^4$ years depending on cloud geometry (Frisch 1994). Prior to that the Sun was immersed in the Local Bubble interior, a region of low density hot plasma ($\sim 0.005 \text{ cm}^{-3}$, 10^6 K) where the heliosphere was probably large (heliopause radius in nose direction, $R_{\text{HP}} \sim 300$ AU, Frisch et al. 2003, FMZL03). An encounter with a diffuse cloud of somewhat higher density, $n(\text{H}^0) \sim 10 \text{ cm}^{-3}$, would contract the heliosphere ($R_{\text{HP}} \sim 15$ AU) with dramatic consequences for galactic cosmic ray modulation, and cosmic ray fluxes at the Earth’s surface. In contrast, today’s heliosphere has $R_{\text{HP}} \sim 100$ AU (see other papers in this issue). Fig. 1 illustrates the motion of the Sun through the Local Bubble and with respect to the Cluster of Local Interstellar Clouds (CLIC).

The influence of the galactic cosmic ray flux on the global electrical circuit (Roble 1991, Tinsley 2000) provides a tentative link between heliosphere dimensions and the 1 AU cosmic ray flux. A positive correlation between low altitude tropospheric cloud cover (≤ 3 km altitude) and neutron monitor counts (a direct tracer of ≥ 0.5 GeV cosmic ray fluxes at Earth, FMZL03) has been established over one solar cycle (~ 1985 – 1995 , Marsh and Svensmark 2000). Evidently cosmic rays interacting with the Earth’s atmosphere produce aerosols which seed water vapor condensation and cloud formation. During the period of time the Sun was immersed in the Local Bubble interior (probably for many millions of years), the heliosphere should have been large with maximum galactic cosmic ray modulation, and in the absence of surrounding ISM neutrals anomalous cosmic rays would have been minimal. If further investigation confirms the correlation between low altitude cloud cover and the 1 AU cosmic ray flux, then it would appear that heliospheric modulation of the galactic and anomalous cosmic ray fluxes has a direct influence on terrestrial climate, and a large heliosphere may be a factor in climate stability.

By analogy, the ISM also sets the boundary conditions for the astrospheres of extrasolar planetary

Fig. 1. ISM within 300 pc of Sun. Figure based on E(B-V) maps of Lucke (1978) and molecular clouds as shown in CO maps of Dame et al. (1987). The three subgroups of the Scorpius-Centaurus Association (SCA) (Upper Scorpius, Upper Centaurus-Lupus, Lower Centaurus-Crux) are shown, as is the location of the Geminga pulsar (towards Orion). The yellow arrow indicates the solar apex motion. The dark blue arrow indicates the bulk flow vector of the CLIC in the LSR. The light blue arrow shows the LIC LSR velocity.

systems (Frisch 1993), although variations in stellar winds must also be considered (Wood et al. 2001). Through the heliosphere-cosmic ray (and by analogy astrosphere-cosmic ray) connection, the physical properties of the present and historical ISM surrounding each system may be a factor in the distribution of stable planetary climates. Although Earth-mass planets have not yet been discovered, over 10% of the more than 100 planets now discovered have semimajor axes near ~ 1 AU (for a list of discovered planets, see URL <http://exoplanets.org>). Spectral types for the central star are typically F, G, or K, and include both giant and main sequence objects. Mass-loss characteristics of cool stars vary with stellar activity levels, which are traced by X-ray surface flux, and decrease with stellar age (Wood et al. 2002). The astrospheres of extrasolar planetary systems may thus vary substantially from the heliosphere, both because of the surrounding ISM and stellar activity levels. These considerations will be important in evaluating the yet-to-be-discovered climates of extrasolar planets, as well as for understanding the climatic history of the Earth.

CHEMICAL COMPOSITION, PHYSICAL PROPERTIES, AND GAS-TO-DUST MASS RATIO OF SURROUNDING CLOUD

The low opacity of the CLIC to the cloud surface in the heliosphere tail direction yields relatively high average ionizations of H° and He° at the solar location. Observations of the ISM towards more distant stars yields sight-line averaged abundances which may be misleading in the presence of

ionization gradients. H^+ is generally unobserved in low column-density sightlines. Both H^0 and H^+ must be included for determining the chemical composition of the LIC, since most of the observed ions are present in both neutral and ionized regions (i.e. He, Ne, Ar, and elements with first ionization potentials less than 13.6 eV). The correct gas-phase abundances are required to evaluate gas-dust mixing and the chemical homogeneity of the ISM.

The LIC is unique in that radiative transfer models are constrained by both observations of the ISM towards nearby stars, and by *in situ* observations of ISM interaction products inside of the solar system (Slavin & Frisch 2002, SF02). The ISM composition and ionization at the solar location has been studied using the radiative characteristics of the material, with the models constrained by observations of ISM towards ϵ CMa (using data of Gry & Jenkins 2001) and by pickup ion data for an assumed $n(H^0)=0.016\pm0.002\text{ cm}^{-3}$ in the LIC. Within the past year new data offers an improved basis for selecting among the 25 models generated by the SF02 study. Studies of O^0/H^0 in the generic ISM now shows that the ratio is relatively constant and ~ 410 (Andre et al. 2002). Voyager observations of anomalous cosmic rays yield data on interstellar argon (Cummings & Stone 2002) which is partially ionized in local ISM ($Ar^0/Ar\sim 0.20-0.24$). More extensive data on pickup ions are also available (Gloeckler & Geiss 2002), as well as better *in situ* observations of He^0 (Witte et al. 1996, and private communication). Two models provide the best match to the range of the available data for an assumed $n(H^0)=0.016\pm0.002\text{ cm}^{-3}$. These two best models are Models 2 and 18, and the parameters for these models are shown in Table 1. These models, based on the assumption that $O/H\sim 400$ PPM in $d<3$ pc ISM (applying results of Andre et al. 2002 to the LISM), show that N and S are present at $\sim 60\%$ of solar abundance while C is mildly subsolar (SF02). For this comparison, solar abundances from Holweber (2001) are assumed. Alternatively, assuming O and C solar abundances from Allende Prieto et al. (2001) would imply unrealistic supersolar C abundances.

The filtration of neutrals in the heliosheath regions is a key unknown quantity. For example, filtration factors for Ar range from ~ 0.64 (Cummings & Stone 2002) to ~ 0.99 (Mueller and Zank 2002). Hydrogen filtration factors are predicted on a purely theoretical basis (e.g. Ripken & Fahr 1983, Izmodenov et al. 1999), however the radiative transfer models also yield an estimate of hydrogen filtration in the heliosheath regions. Gloeckler & Geiss (2002) derive the filtration factor for H^0 by comparing H/He measured in the pickup ion population with H^0/He^0 for the nearby ISM. Using their approach, Frisch and Slavin (2002) have shown that Models 2 and 18 predict H filtration factors (~ 0.43) which yield agreement between pickup ion $n(H^0)$ values and the model predictions for the solar location. This is illustrated in Fig. 2.

The gas-to-dust mass ratio ($R_{g/d}$) may be calculated from the radiative transfer models, and the results suggest that interstellar gas and dust are not well mixed in the LIC (Frisch & Slavin 2002). When missing mass arguments and an assumed solar reference abundance are applied to Model 18, a gas-to-dust mass ratio of ~ 180 is found; subsolar ISM reference abundances yield values a factor of ~ 3 larger. However, *in situ* dust observations of interstellar dust by Ulysses, Galileo, and Cassini yield a gas-to-dust mass ratio of $\sim 125\pm 18$ (Frisch et al. 1999; Landgraf et al. 2000; Frisch & Slavin 2002). One explanation for the difference between these two values for $R_{g/d}$ would be poor gas-dust mixing as a superbubble shell fragment sweeps up both gas and dust in the ISM. Most of the mass of the interstellar dust grains detected within the solar system is carried by grains with masses $> 10^{-13}$ g (or radius $> 0.2\text{ }\mu\text{m}$). The gyro-radius for a grain with mass 10^{-13} g (radius $\sim 0.2\text{ }\mu\text{m}$) in a weak magnetic field ($\sim 3\text{ }\mu\text{G}$) is ~ 0.3 pc (Gruen & Landgraf 2000), which is greater than the distance to the LIC edge in the upstream direction. Thus massive dust grains decouple from gas in the upstream LIC, and the $R_{g/d}$ over the small scales sampled by the *in situ* data differs from the value averaged over a typical long interstellar sightlines. The result is that $R_{g/d}$ may vary over sub-parsec length scales. Other explanations, such as poorly understood ISM abundances, are also possible.

Table 1. LIC Parameters for Best Models^a

Quantity	Model 2	Model 18
<u>Assumed Parameters:</u>		
n_{tot} (cm ⁻³)	0.273	0.300
$\log T_{\text{h}}$ (K)	6.0	6.1
B_{o} (μG)	5.0	3.0
$N_{\text{H}^{\text{o}}}$ (10^{17} cm ⁻²) ^b	6.5	6.5
<u>Results for ISM at Solar Location:</u>		
$n(\text{H}^{\text{o}})$ (cm ⁻³)	0.208	0.242
$n(\text{He}^{\text{o}})$ (cm ⁻³)	0.015	0.017
$n(\text{e})$ (cm ⁻³)	0.098	0.089
$\chi(\text{H})^{\text{c}}$	0.287	0.234
$\chi(\text{He})^{\text{c}}$	0.471	0.448
$\log N_{\text{tot}}$ (cm ⁻²) ^b	18.03	17.98
$N(\text{H}^{\text{o}})/N(\text{He}^{\text{o}})^{\text{b}}$	11.6	12.1
T (K)	8,230	8,140

^a Models are from Slavin & Frisch (2002) and Frisch & Slavin (2003). $n_{\text{tot}} = n(\text{H}^{\text{o}}) + n(\text{H}^+)$, while $\log N_{\text{tot}} = N(\text{H}^{\text{o}}) + N(\text{H}^+)$. ^b The column density $N(\text{H}^{\text{o}})$ is the H^{o} column density between the Sun and cloud surface, etc. In this case the cloud is the sum of the two interstellar cloudlets within ~ 3 pc of the Sun in the ϵ CMa direction (Gry & Jenkins 2001). ^c $\chi(\text{H}, \text{He})$ is the ionization fraction of element H, He.

SOLAR MOTION AND ORIENTATION WITH RESPECT TO THE INTERSTELLAR MAGNETIC FIELD

The Sun moves through space and has recently emerged from the interior of the Local Bubble and entered an outflow of ISM from the Scorpius-Centaurus Association (Fig. 1). Hipparcos results yield a solar motion in the local standard of rest of ~ 13.4 km s⁻¹ towards the galactic coordinates $l=28^\circ$, $b=+32^\circ$ (Dehnen & Binney 1998).

The north ecliptic pole is directed towards $l=96^\circ$, $b=30^\circ$, giving an ecliptic plane tilted by $\sim 60^\circ$ with respect to the galactic plane. Observations of the interstellar magnetic field directly outside of the heliosphere have been elusive, but evidence for a weak magnetic field ($< 3 \mu\text{G}$) is given by starlight polarization caused by interstellar dust grains aligned by nearby magnetic fields (Tinbergen 1982, Frisch 1990). The patch of ISM causing this polarization is located towards $l=0 \pm 20^\circ$, $b=-20 \pm 25^\circ$, and is close to the ecliptic plane towards the nose direction of the heliosphere (see Fig. 3). The polarization data show a nearby magnetic field orientation which is approximately parallel to the galactic plane and directed towards $l \sim 70^\circ$. The classical interstellar dust grains which cause the polarization are charged, and pile up in the heliosheath regions as they are deflected around the heliosphere (Frisch et al. 1999). Although the region of maximum polarization closely follows the ecliptic plane, it also coincides with the CLIC LSR upstream direction (Fig. 3). Observations by Voyager 1 and 2 observations of ~ 3 kHz emission in the outer heliosphere also indicate the interstellar magnetic field direction is parallel to the galactic plane (Kurth & Gurnett 2003). The combined ecliptic tilt and interstellar magnetic field direction produce a north/south asymmetry in




Fig. 2. Comparison between H^0 in the pickup ion population and model predictions for $n(H^0)$ for the 25 models of Slavin & Frisch (2002). The ordinate in each plot is the interstellar $n(H^0)$ value that results after the pickup ion measurement of $n(H^0)$ at the termination shock is corrected for heliosheath filtration (F) using H^0/He^0 at the solar location from the Slavin & Frisch (2002) for each model. The abscissa for the plot is the predicted $n(H^0)$ value at solar location. The viable models are shown with pickup ion data uncertainties plotted, and the three best models based on Mg^+/Mg^0 and C^+/C^{+*} ratios towards ϵ CMa have boxes around the data points. The dotted line illustrates perfect agreement between models and data. Models 2 and 18 appear to provide the best match to the combined pickup ion and ϵ CMa data when $n(He^0)=0.016\pm0.002\text{ cm}^{-3}$ in the LIC. The points for Models 16 and 17, which differ only in the FUV radiation field, are superimposed on each other. (Adopted from Frisch & Slavin 2002.)

the heliosphere configuration which is visualized in Fig. 4.

KINEMATICS AND TURBULENCE OF NEARBY ISM

The LIC is one of a group of cloudlets which have a bulk flow motion in the Local Standard of Rest (LSR) of $-17\pm5\text{ km s}^{-1}$, from the upstream direction $l\sim2.3^\circ$, $b\sim-5.2^\circ$ (Frisch et al. 2002). This group of cloudlets has been referred to as the Cluster of Local Interstellar Clouds, or CLIC. The dispersion of the cloudlets about the bulk flow represents turbulence in the nearby ISM. Models 2 and 18 indicate that $n(H^0)=0.208\text{--}0.242\text{ cm}^{-3}$ in the LIC. The hydrogen column density towards the nearest star, α Cen (1.3 pc), has been reported as $N(H^0)=0.41\text{--}1.04\text{ }10^{18}\text{ cm}^{-2}$ (Linsky & Wood 1996), yielding a filling factor of $f=0.5\text{--}1$ for ISM in this sightline. Two cloudlets near the Sun have velocities consistent with the absorption component towards α Cen, the “G” cloud (Lallement & Bertin 1992) and the Apex Cloud seen towards Aquila and Ophiuchus (Frisch 2003). The LSR upstream direction velocity vectors for the G and Apex Clouds are, respectively, $(V, l, b)=(-17.7\text{ km s}^{-1}, 350.5^\circ, 9.2^\circ)$ and $(-23.4\text{ km s}^{-1}, 4.8^\circ, 2.1^\circ)$. For comparison the LSR upstream LIC motion is $(-15.8\text{ km s}^{-1}, 346.1^\circ, 0.2^\circ)$. The Sun is located within a few parsecs of the downstream edge of the bulk flow. The smaller velocity of the LIC, located at the leading edge of the CLIC bulk flow, suggests flow deceleration, while the dispersion of the CLIC velocities reflects turbulence. These flow parameters are derived from observations of 96 interstellar absorption line components (mainly Ca^+ data) towards 60 stars which sample the ISM within 30 parsecs of the Sun (Frisch, Grodnicki & Welty 2002). The LSR upstream

direction of the CLIC bulk flow is within $\sim 20^\circ$ of the observed tangent point of Loop I, which has been formed by successive epochs of star formation in the Scorpius-Centaurus Association over the past 10^7 years. The CLIC gas has been attributed to a superbubble shell fragment associated with this star formation, with possible implications for the chemical abundance pattern for this nearest gas. For such an origin, the CLIC would be composed of swept up material consisting of ejecta from previous epochs of star formation in the SCA, the ambient ISM, and recently ejected nucleosynthetic products. This scenario provides a hybrid method for the dispersal of nucleosynthetic products – e.g. cloud collisions and turbulence within the outflow.




Fig. 3. Plot of an indicator of the nearby interstellar magnetic field, in galactic coordinates. The bars show the direction of the electric vector polarization, which is parallel to the interstellar magnetic field direction, for several nearby stars (Tinbergen 1982). The arrow shows the likely direction of the interstellar magnetic field near the Sun based on these data. The curved line shows the ecliptic plane. The region of maximum polarization follows the ecliptic plane, but is also in the LSR upstream CLIC direction (box). A star is plotted at the heliosphere nose direction in heliocentric coordinates. The classical interstellar dust grains which polarize optical radiation pile up in the heliosheath regions as they are deflected around the heliosphere (Frisch et al. 1999). (Figure adopted from Frisch & Slavin 2002.)

CONCLUSIONS

Why study nearby ISM? Taking an anthropomorphic viewpoint, the ISM dominates the interplanetary medium throughout most of the heliosphere. Heliosphere models show that heliosphere properties, and by analogy exoplanetary system astrospheres, are dramatically affected by the physical properties of the surrounding interstellar cloud. From an astronomical viewpoint, the $d < 30$ pc ISM is the best region for studying the physical processes which operate in the ISM because of the unusual range of data available for constraining radiative transfer models. These data include the ISM properties within the solar system, which trace the neutral component. Accurate ISM models permit basic axioms underlying ISM studies to be tested, such as the assumption that both gas and dust must be included when evaluating the chemical composition of the ISM.

Fig. 4. Visualization of the asymmetric heliosphere caused by the apparent tilt of the ecliptic plane with respect to the local interstellar magnetic field (from Hanson et al. 2002). The heliosphere morphology is based on an MHD simulation by Linde et al. (1998), which includes the relative orientation and ram pressures of the interstellar and solar wind magnetic fields. Note the constellation of Orion is viewed through the heliosphere tail. The movie containing this visualization can be downloaded from: <http://cs.indiana.edu/~soljourn>. (Used with permission.)

ACKNOWLEDGEMENTS

The author would like to gratefully acknowledge research support from NASA grants NAG5-6405, and NAG5-11005, and NAG5-8163.

REFERENCES

- Adams, T. F., and P. C. Frisch, High-resolution observations of the Lyman alpha sky background, *Astrophys. J.*, **212**, 300–308, 1977.
- Allende Prieto, C., D. L. Lambert, and M. Asplund, The forbidden abundance of oxygen in the Sun, *Astrophys. J. Lett.*, **556**, L63–L66, 2001.
- Andre, M., et al., Oxygen gas phase abundance revisited, submitted to *Astrophys. J.*, 2002.
- Cummings, A. C., E. C. Stone, and C. D. Steenberg, Composition of anomalous cosmic rays and other heliospheric ions, *Astrophys. J.*, **578**, 194–210, 2002.
- Dame, T. M., H. Ungerechts, R. S. Cohen, E. J. de Geus, I. A. Grenier, J. May, D. C. Murphy, L. A. Nyman, and P. Thaddeus, A composite CO survey of the entire Milky Way, *Astrophys. J.*, **322**, 706–720, 1987.
- Dehnen, W., and J. J. Binney, Local stellar kinematics from Hipparcos data, *MNRAS*, **298**, 387–394, 1998.
- Frisch, P. C., Characteristics of the local interstellar medium, in *Physics of the Outer Heliosphere*, edited by S. Grzedzielski and D. E. Page, pp. 19–22, Pergamon Press, Oxford, 1990.
- Frisch, P. C., G-star astropauses - A test for interstellar pressure, *Astrophys. J.*, **407**, 198–206, 1993.
- Frisch, P. C., Morphology and ionization of the interstellar cloud surrounding the solar system, *Science*, **265**, 1423–1426, 1994.
- Frisch, P. C., Journey of the Sun, <http://xxx.lanl.gov/>, astro-ph/9705231, 1997.
- Frisch, P. C., and J. D. Slavin, Chemical composition and gas-to-dust mass ratio of the nearest interstellar matter, submitted to *Astrophys. J.*, , 2002.
- Frisch, P. C., L. Grodnicki, and D. E. Welty, The velocity distribution of the nearest interstellar gas, *Astrophys. J.*, **574**, 834–846, 2002.
- Frisch, P. C., H. R. Mueller, G. P. Zank, and C. Lopate, *Galactic environment of the Sun and Stars: Interstellar and interplanetary material*, in press, Cambridge: Cambridge University Press, 2003.
- Frisch, P. C., Dorschner, J. M. and Geiss, J., et al., Dust in the Local Interstellar Wind, *Astrophys. J.*, **525**, 492–516, 1999.
- Gloeckler, G., and G. Geiss, Composition of the local interstellar medium as diagnosed with pickup ions, *Adv. Space Res.*, this issue, 2003.
- Gruen, E., and M. Landgraf, Collisional consequences of big interstellar grains, *J. Geophys. Res.*, **105**, 10291–10298, 2000.
- Gry, C., and E. B. Jenkins, Local clouds: Ionization, temperatures, electron densities and interfaces, from GHRS and IMAPS spectra of epsilon Canis Majoris, *A&A*, **367**, 617–628, 2001.
- Hanson, A. J., C.-W. Fu, E. A. Wernert, and P. C. Frisch, Case study: Constructing the solar journey, *Preprint*, 2002.
- Holweger, H., Photospheric abundances: Problems, updates, implications, <http://xxx.lanl.gov/>, astro-ph/0107426, 2001.

- Izmodenov, V. V., J. Geiss, R. Lallement, G. Gloeckler, V. B. Baranov, and Y. G. Malama, Filtration of interstellar hydrogen in the two-shock heliospheric interface: Inferences on the local interstellar cloud electron density, *J. Geophys. Res.*, **104**, 4731–4742, 1999.
- Kurth, W. S., and D. A. Gurnett, On the source location of low-frequency heliospheric radio emissions, submitted to *J. Geophys. Res.*, 2003.
- Lallement, R., and P. Bertin, Northern-Hemisphere observations of nearby interstellar gas - Possible detection of the local cloud, *A&A*, **266**, 479–485, 1992.
- Landgraf, M., W. J. Baggaley, E. Grün, H. Krüger, and G. Linkert, Aspects of the mass distribution of interstellar dust grains in the solar system from in situ measurements, *J. Geophys. Res.*, **105**, 10343–10352, 2000.
- Linde, T. J., A three-dimensional adaptive multifluid MHD model of the heliosphere, Ph.D. thesis, Univ. of Michigan, Ann Arbor, 1998, small <http://hpcc.engin.umich.edu/CFD/publications>.
- Linde, T. J., T. I. Gombosi, P. L. Roe, K. G. Powell, and D. L. DeZeeuw, Heliosphere in the magnetized local interstellar medium: Results of a three-dimensional MHD simulation, *J. Geophys. Res.*, **103**, 1889–1904, 1998.
- Linsky, J. L., and B. E. Wood, The alpha Centauri line of sight: D/H ratio, physical properties of local interstellar gas, and measurement of heated hydrogen (the ‘hydrogen wall’) near the heliopause, *Astrophys. J.*, **463**, 254–270, 1996.
- Lucke, P. B., The distribution of color excesses and interstellar reddening material in the solar neighborhood, *A&A*, **64**, 367–377, 1978.
- Müller, H., and G. P. Zank, Modeling heavy ions and atoms throughout the heliosphere, in press, “*Solar Wind 10; AIP Conference Proceedings*”, 2002.
- Marsh, N. D., and H. Svensmark, Low cloud properties influenced by cosmic rays, *Physical Review Letters*, **85**, 5004–5007, 2000.
- Mueller, H. R., G. P. Zank, and P. C. Frisch, Effect of different possible interstellar environments on the heliosphere: A numerical study, in “*The Outer Heliosphere: The Next Frontiers*”, edited by K. Scherer, COSPAR Colloquium Series, 11, 329–333, 2001.
- Mueller, H. R., G. P. Zank, and P. C. Frisch, Heliospheric response to different possible interstellar environments, *in preparation*, 2002.
- P. C. Frisch, Local interstellar matter: The Apex Cloud, submitted to *Astrophys. J.*, 2003.
- Ripken, H. W., and H. J. Fahr, Modification of the local interstellar gas properties in the heliospheric interface, *A&A*, **122**, 181–192, 1983.
- Roble, R. G., On modeling component processes in the earth’s global electric circuit, *Journal of Atmospheric and Terrestrial Physics*, **53**, 831–847, 1991.
- Slavin, J. D., and P. C. Frisch, The Ionization of Nearby Interstellar Gas, *Astrophys. J.*, **565**, 364–379, 2002.
- Tinbergen, J., Interstellar polarization in the immediate solar neighborhood, *A&A*, **105**, 53–64, 1982.

- Tinsley, B. A., Influence of solar wind on the global electric circuit, and inferred effects on cloud microphysics, temperature, and dynamics in the Troposphere, *Space Science Reviews*, **94**, 231–258, 2000.
- Witte, M., M. Banaszkiewicz, and H. Rosenbauer, Recent results on the parameters of the interstellar helium from the ULYSSES/GAS experiment, *Space Sci. Rev.*, **78**, 289–2396, 1996.
- Wood, B., J. Linsky, H. Müller, and G. Zank, Observational estimates for the mass-loss rates of alpha; Centauri and Proxima Centauri using Hubble Space Telescope Ly-alpha spectra, *Astrophys. J. Lett.*, *547*, L49–L52, 2001.
- Wood, B., H. Müller, G. Zank, and J. Linsky, Measured mass-loss rates of solar-like stars as a function of age and activity, *Astrophys. J. Lett.*, **574**, 412–425, 2002.
- Zank, G. P., and P. C. Frisch, Consequences of a change in the Galactic environment of the Sun, *Astrophys. J.*, **518**, 965–973, 1999.

Manuscript received 3 *December* 2002, revised 13 *February* 2003, accepted 13 *February* 2003.

This figure "berry.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0302254v2>

This figure "fig4.jpg" is available in "jpg" format from:

<http://arXiv.org/ps/astro-ph/0302254v2>